



Actero™ Buffered Peptone Water Product Information

Catalogue No.	Description
FCM-066	Actero™ Buffered Peptone Water (500 G)
FCM-065	Actero™ Buffered Peptone Water (2 KG)
FCM-064	Actero™ Buffered Peptone Water (10 KG)
FCM-191	Actero™ Buffered Peptone Water MediaPouch 10L / 200g
FCM-193	Actero™ Buffered Peptone Water 20L / 400g

INTENDED USE

Buffered Peptone Water is used for pre-enriching damaged *Salmonella* spp. from various food sources. Edel and Kampelmacher¹ noted that food preservation techniques such as heat, desiccation, preservatives, high osmotic pressures or pH changes can cause sub-lethal injury to *Salmonella* Microorganisms. Pre-enrichment with Buffered Peptone Water allows for repair of cell damage by maintaining a high pH over the pre-enrichment period. The high pH capacity is especially useful for vegetable samples which have a low buffering capacity.

Formula* per Liter:

Pancreatic Digest of Gelatin.....10.0g
Disodium Phosphate.....3.5g
Sodium Chloride.....5.0g
Monopotassium Phosphate.....1.5g

Final pH: 7.2 ± 0.2 at 25°C

** Grams per liter may be adjusted or formula supplemented to obtain desired performance.*

Storage Instructions:

On receipt, store the sealed bottle containing the dehydrated medium at 2 to 30°C. Once opened and recapped, place the container in a low humidity environment at the same storage temperature. Protect it from moisture and light. The dehydrated medium should be discarded if it is not free flowing or if the color has changed from the original light beige.

QUALITY CONTROL SPECIFICATIONS

1. The powder is homogeneous, free flowing and light beige.
2. Visually the prepared medium is clear and pale yellow.
3. Expected cultural response after 18-24 hours at 35°C.

Organism	Result
<i>Escherichia coli</i> ATCC 25922	Growth
<i>Salmonella enteritidis</i> ATCC 13076	Growth
<i>Salmonella typhimurium</i> ATCC 14028	Growth

PREPARATION

Mix 20 grams of the medium in one liter of purified water until evenly dispersed. Heat with repeated stirring and boil for one minute to dissolve completely. Distribute and autoclave at 121°C for 15 minutes.

References

1. Edel, W., and E. H. Kampelmacher. 1973. Bull. World Health. Org. 48:167-174.



