



Actero™ Baird Parker Agar Product Information

Catalogue No.	Description
FCM-060	Actero™ Baird Parker Agar (500 G)
FCM-059	Actero™ Baird Parker Agar (2 KG)
FCM-058	Actero™ Baird Parker Agar (10 KG)

INTENDED USE

Baird Parker Agar is used with Egg Yolk Tellurite Enrichment for the detection, isolation and enumeration of coagulase positive *staphylococci* in foods and cosmetics. Potassium Tellurite and Lithium Chloride selectively suppress the growth of most bacteria but allow the growth of Staphylococci which is also enhanced by the addition of sodium pyruvate and glycine. The differentiation of coagulase-positive *staphylococci* is noted by the reduction of Potassium Tellurite which results in a black colony. *Staphylococci* that contain lecithinase break down the egg yolk and cause a clear zone. Opaque zones may form as a result of lipase activity.

Formula* per Liter:

Casein Digest of Peptone 10.0g
Sodium Pyruvate 10.0g
Glycine 12.0g
Beef Extract..... 5.0g
Yeast Extract 1.0g
Lithium Chloride..... 5.0g
Agar..... 17.0g
Final pH: 7.0 ± 0.2 at 25°C

* Grams per liter may be adjusted or formula supplemented to obtain desired performance.

PREPARATION

Mix 60 grams of the medium in one liter of purified water until evenly dispersed. Heat with repeated stirring and boil for one minute to dissolve completely. Distribute and autoclave at 121°C for 15 minutes. After cooling to 45-50.0°C add 10 mL of a sterile 1% Tellurite Supplement and 50 mL of Egg Yolk Emulsion. Mix thoroughly, dispense as desired.

QUALITY CONTROL SPECIFICATIONS

1. The powder is homogeneous, free flowing, and beige.
2. Visually the prepared medium is clear to slightly hazy and light amber. The prepared enriched medium is canary yellow and opaque.
3. Expected cultural response after 24-48 hours at 35°C.

Organism	Growth	Colony Color	Reactions
<i>Escherichia coli</i> ATCC® 25922	Inhibition	-	N/A
<i>Proteus mirabilis</i> ATCC® 12453	Good	Black colonies w/o halo	Lipase & Lecithinase -
<i>Bacillus subtilis</i> ATCC® 6633	Inhibition	-	N/A
<i>Staphylococcus aureus</i> ATCC® 25923	Excellent	Black colonies w/ halo	Lipase & Lecithinase +
<i>Staphylococcus epidermidis</i> ATCC® 12228	Good	Grey/Black colonies w/o	Lipase & Lecithinase -

STORAGE INSTRUCTIONS:

On receipt, store the sealed bottle containing the dehydrated medium at 2 to 30°C. Once opened and recapped, place the container in a low humidity environment at the same storage temperature. Protect it from moisture and light. The dehydrated medium should be discarded if it is not free flowing or if the color has changed from the original beige or color.

